

Evidence for h_c production from ψ' at CLEO

Amiran Tomaradze


(Northwestern University)

Representing the CLEO Collaboration

We present the results of our search for the $h_c(^1P_1)$ state of charmonium in reaction

$$e^+e^- \rightarrow \psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c)$$

using CLEO data.



This Presentation Includes:

- Importance of Identifying h_c
- Prior Experimental Searches for h_c
- CLEO Searches and Results

THE IMPORTANCE OF $h_c(^1P_1)$

- Of the 8 bound states of charmonium, $h_c(^1P_1)$ is the only one ‘undiscovered’ (or unconfirmed) in the last 30 years since the discovery of J/ψ .
- There is an important physics reason to identify h_c .
 - The spin-independent part of the QCD $q\bar{q}$ interaction is well-established.
 - The spin-dependent part is not.

In particular, the $\vec{s}_1 \cdot \vec{s}_2$ spin–spin, or **hyperfine interaction is not well understood**, because there is little experimental data to provide the required constraints for theory.
- The primary experimental data required for understanding the $q\bar{q}$ hyperfine interaction is hyperfine, or spin-singlet/spin-triplet splitting:

$$\Delta M_{hf}(nL) \equiv \langle M(n^3L_J) \rangle - M(n^1L_{J=L})$$

- No spin-singlet states have so far been identified in bottomonium. **Charmonium is our only source.**

THE IMPORTANCE OF $h_c(^1P_1)$

- For nearly 20 years, the only hyperfine splitting known was that for the $1S$ states of charmonium

$$\Delta M_{hf}(1S) = M(J/\psi) - M(\eta_c) = 116 \pm 2 \text{ MeV}$$

- Very recently, Belle, CLEO and BaBar, succeeded in identifying η'_c , with the rather surprising result that

$$\Delta M_{hf}(2S) = M(\psi') - M(\eta'_c) = 48 \pm 4 \text{ MeV}$$

Potential model and quenched lattice calculations predicted larger $\Delta M_{hf}(2S)$.

- It is of great importance to find out how the hyperfine interaction manifests itself in P states, i.e., to find $\Delta M_{hf}(1P) \equiv M(\langle ^3 P_J \rangle) - M(^1P_1)$
- With scalar confinement, $\Delta M_{hf}(1P) = 0$ is expected. It is necessary to determine if this is true.
- The c.o.g. of 3P states, $M(\langle ^3 P_J \rangle)$, is well measured, $M(\langle ^3 P_J \rangle) = 3525.3 \pm 0.1 \text{ MeV}$,

What we need is to identify h_c and make a precision measurement of its mass.

$h_c(^1P_1)$ - EXPERIMENTAL

- 1982: Crystal Ball (SLAC), not observed in $M = 3440 - 3543$ MeV

$$\psi' \rightarrow \pi^0 h_c, \quad h_c \rightarrow \gamma \eta_c, \quad B_{in} \times B_{out} < 0.32 \times 10^{-2}$$

NO

- 1992: E760 (FNAL), $p\bar{p}$ in $M = 3523 - 3527$ MeV, $\mathcal{L} = 16 \text{ pb}^{-1}$

$$p\bar{p} \rightarrow (h_c) \rightarrow \pi^0 J/\psi, \quad B_{in} \times B_{out} \approx 2 \times 10^{-7}, \quad \sim 3\sigma$$

$$M = 3526.2 \pm 0.3 \text{ MeV}, \quad \Gamma \leq 1.1 \text{ MeV}$$

YES

- 2004: E835 (FNAL), $p\bar{p}$ in $M = 3523 - 3529$ MeV, $\mathcal{L} = 45 \text{ pb}^{-1}$

$$p\bar{p} \rightarrow (h_c) \rightarrow \pi^0 J/\psi, \quad B_{in} \times B_{out} \leq 1 \times 10^{-7} \text{ (90\%CL)}$$

NO

$$p\bar{p} \rightarrow (h_c) \rightarrow \eta_c \gamma, \quad M = 3525.8 \pm 0.2(\text{stat}) \pm 0.2(\text{syst}) \text{ MeV}, \quad \sim 3\sigma$$

PRELIMINARY YES

- 2004: Belle and BaBar from B decays

PRELIMINARY NO

CLEO Searches and Results


We analyze $\sim 6pb^{-1}$ of CLEO III and CLEO-c data with estimated $\sim 3.0 \times 10^6$ ψ' events, for $\psi' \rightarrow \pi^0 h_c$, $\pi^0 \rightarrow \gamma\gamma$, $h_c \rightarrow \gamma\eta_c$.

1. We search for this channel without using η_c decays (INCLUSIVE approach).
2. We search for this channel using six dominant η_c decay modes (EXCLUSIVE approach).

In both methods we search for h_c in the mass recoiling against π^0 from decay $\psi' \rightarrow \pi^0 h_c$.

This method benefits from the excellent resolution of the CLEO calorimeter.

ALL RESULTS ARE PRELIMINARY !



CLEO-c Calorimeter

- 7800 CsI(Tl) crystals
- 93% of 4π coverage
- Energy resolution: $\sigma_E/E = 1.5\%$ at 5 GeV, 4% at 100 MeV.

Inclusive Analyses, Event Selection

Two independent analyses have been done. The results from the two are consistent.

I will describe one of them in detail, and will later mention the differences between the two analyses.

We use the following selection criteria to select

$\psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c)$ events.

- $N_{shower} \geq 3$.
- $N_{track} \geq 2$.

Photon shower selection

Standard photon selections were used.

- $E_\gamma > 30$ MeV, if shower is in the barrel
- $E_\gamma > 50$ MeV, if shower is in the endcap

Track selection

The selection of charged particles which is used for rejection of $(\psi' \rightarrow \pi^+\pi^- J/\psi)$ and $(J/\psi \rightarrow \text{charged})$ events is done using the standard quality cuts.

Event Selection

- Reconstruction of π^0 's: $M_{\gamma\gamma}=135\pm 15$ MeV.


The fitted π^0 's are used. We require that there be only one π^0 in the event with recoil mass in range of expected h_c mass of 3526 ± 30 MeV.

- Reject $\psi' \rightarrow \pi^+\pi^- J/\psi$ events by cutting out recoils against $\pi^+\pi^-$.
- Reject $\psi' \rightarrow \pi^0\pi^0 J/\psi$ events by cutting out recoils against $\pi^0\pi^0$.
- Reject event if mass of all charged particles is around invariant mass of J/ψ .
- Reject candidate hard γ (from $h_c \rightarrow \gamma\eta_c$ decays) which makes π^0 or η with any other γ 's.
- Cut on hard γ energy, $E_\gamma=503\pm 40$ MeV, which corresponds to $M(h_c)=3526\pm 47$ MeV.

Monte Carlo Studies

- We analyzed a sample of $\sim 12 \times 10^6$ generic ψ' Monte Carlo events (events containing all measured ψ' decays except those via h_c) in four separate samples, each with approximately the same size ($\sim 3 \times 10^6$) as the data.
- The 10,000 signal MC events for the channel $\psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c)$ were simulated, assuming:
 $M(h_c) = 3526 \text{ MeV}, \Gamma(h_c) = 0.0, 0.5, 0.9 \text{ MeV}$
 $M(\eta_c) = 2982 \text{ MeV}, \Gamma(\eta_c) = 24.8 \text{ MeV}.$
- The signal MC events were added in to the generic MC assuming:
 $B(\psi' \rightarrow \pi^0 h_c) \times B(h_c \rightarrow \gamma\eta_c) = 5.0 \times 10^{-4}.$
- The event selection criteria applied to the Monte–Carlo samples were identical to those applied to the data.

Recoil Mass Resolution at h_c



Recoil mass distribution against π^0 in reaction $\psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c)$ in signal Monte-Carlo, for input $M(h_c)=3526$ MeV, $\Gamma(h_c)=0$ MeV.

The full curve shows fit with a double Gaussian ($\sigma_1=1.3$ MeV, $\sigma_2=3.7$ MeV, $\text{area}_2/\text{area}=0.43$) which gives the efficiency=16%. These parameters are used to fit the signal in the data.

Background Shape

The **background** in the data has been fitted in three ways:

- ARGUS shape,





$$y = x \times \sqrt{1 - (x/a)^2} \times \exp(b \times [1 - (x/a)^2]),$$

- 3 parameter polynomial shape,
- Background shape from Monte–Carlo.

Significance levels are obtained as


$\sigma \equiv \sqrt{-2 \ln(L_0/L_{max})}$, where L_{max} and L_0 are the likelihoods of the fits with and without the h_c resonance.

GEANT MC, π^0 recoil mass in GeV



	$M(h_c)$ (MeV)	$B_1 \times B_2$ (10^4)	signif.
input	3526.0	5.0	–
1st sample	3525.9 ± 0.6	4.5 ± 1.0	4.5σ
2nd sample	3526.3 ± 0.6	4.6 ± 1.0	4.3σ
3rd sample	3525.2 ± 0.4	5.5 ± 1.1	5.3σ
4th sample	3525.7 ± 0.5	3.7 ± 1.1	3.4σ
$\chi^2/\text{d.o.f.}$	1.4	1.0	–
Sum	3525.6 ± 0.3	4.4 ± 0.6	8.4σ

CLEO DATA



Distribution of the recoiling mass against π^0 in data.
(Double Gaussian + ARGUS background)

The best fit results are

$$N(h_c) = 156 \pm 48,$$

$$\text{significance}(h_c) = 3.3\sigma,$$

$$M(h_c) = 3524.4 \pm 0.7 \text{ MeV}.$$

Alternative Inclusive Analysis

- Independent inclusive search for h_c
- Different but overlapping event selection
- Select candidates based on recoil against $\pi^0\gamma$ (η_c mass) rather than γ energy
- Consistent results with previously described analysis

Thus our PRELIMINARY CLEO results from two inclusive analyses are:

- $M(h_c) = 3524.8 \pm 0.7(\text{stat}) \pm \sim 1(\text{syst}) \text{ MeV}$.
- $B(\psi' \rightarrow \pi^0 h_c) \times B(h_c \rightarrow \gamma \eta_c) = (2-6) \times 10^{-4}$,
- The significance of h_c detection $> 3 \sigma$.

Systematics in $M(h_c)$

Estimates of systematic errors in $M(h_c)$ have been made by studying the following:

- π^0 energy scale
- background shapes
- Monte Carlo input/output differences
- non-resonant background
- assumed h_c width
- binning effects
- cut variations

and finally

- the difference in $M(h_c)$ in the two inclusive analyses

These studies also convinced us that the significance of the h_c signal is safely $>3 \sigma$.

I now want to present the preliminary results of an exclusive analysis.

Exclusive Analysis

$$\psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c), \eta_c \rightarrow \text{hadrons}$$

The six η_c decay modes which have reasonably high PDG04 branching ratios have been studied:

$$\eta_c \rightarrow K_s K^\pm \pi^\mp \quad \text{BR}=(1.8\pm 0.6)\%$$

$$\eta_c \rightarrow K^+ K^- \pi^0 \quad \text{BR}=(0.9\pm 0.3)\%$$

$$\eta_c \rightarrow K^+ K^- \pi^+ \pi^- \quad \text{BR}=(2.0\pm 0.7)\%$$

$$\eta_c \rightarrow 2\pi^+ 2\pi^- \quad \text{BR}=(1.2\pm 0.4)\%$$

$$\eta_c \rightarrow \pi^+ \pi^- \eta, \quad \eta \rightarrow \gamma\gamma \quad \text{BR}=(1.3\pm 0.4)\%$$


$$\eta_c \rightarrow \pi^+ \pi^- \eta, \quad \eta \rightarrow \pi^+ \pi^- \pi^0 \quad \text{BR}=(0.7\pm 0.2)\%$$

Event Selection

- Standard photon and track selection criteria were used.
- π^0 : fitted with mass pull $< 3\sigma$, $E_\gamma > 30$ MeV.
- η 's were detected in two decay modes:
 - $\eta \rightarrow \gamma\gamma$: mass pull $< 3\sigma$
 - $\eta \rightarrow \pi^+\pi^-\pi^0$: invariant mass within 20 MeV of nominal.
- $\psi' \rightarrow \pi^+\pi^-J/\psi$ rejected by cutting out recoils against $\pi^+\pi^-$.
- K_S : flight significance $> 3\sigma$, invariant mass of $\pi^+\pi^-$ within 10 MeV of nominal.
- The charged pions and kaons have been identified by RICH and dE/dx .
- The total energy–momentum conservation of the event has been required.
- The invariant mass of the η_c candidates are required to be close to the nominal η_c mass (within 50 MeV).

CLEO DATA

$$\Psi' \rightarrow \pi^0 h_c \rightarrow (\gamma\gamma)(\gamma\eta_c), \eta_c \rightarrow \text{hadrons}$$




A Useful Comparison.


$\psi' \rightarrow \pi^0 h_c \rightarrow \pi^0 \gamma \eta_c$ versus $\psi' \rightarrow \gamma \eta_c$

Decay Mode	$N(\pi^0 \gamma \eta_c)$	$N(\gamma \eta_c)$, normalized to $N(\pi^0 \gamma \eta_c)$
$K^+ K^- \pi^+ \pi^-$	4	1.6 ± 0.9
$\pi^+ \pi^- \pi^+ \pi^-$	8	5.5 ± 1.5
$K_S K \pi$	2	4.0 ± 0.9
$K^+ K^- \pi^0$	1	2.0 ± 0.8
$\pi^+ \pi^- \eta(\gamma\gamma)$	0	2.0 ± 0.4
$\pi^+ \pi^- \eta(\pi^+ \pi^- \pi^0)$	1	1.0 ± 0.5

The observed counts in the h_c decay to η_c are in agreement with what is expected for the direct decay of ψ' to η_c .



CLEO DATA



The results are

$$N(h_c) = 15.0 \pm 4.2,$$

$$\text{significance}(h_c) \sim 5 \sigma,$$

$$M(h_c) = 3524.4 \pm 0.9(\text{stat}) \text{ MeV}.$$

Note that the significance is calculated using likelihood differences.

The background estimation by using η_c sidebands, or by using Monte–Carlo events, yielded consistent results.

No estimate of the systematic uncertainty in $M(h_c)$ has been made so far.

SUMMARY

- We have analyzed $\sim 3.0 \times 10^6$ ψ' from CLEO III and CLEO-c to search for $h_c(^1P_1)$ production in the reaction $\psi' \rightarrow \pi^0 h_c$, $h_c \rightarrow \gamma \eta_c$ by two methods
 1. INCLUSIVE – which does not use η_c decay modes,
 2. EXCLUSIVE – which uses six hadronic decay modes of η_c .

- In the recoil mass spectrum of π^0 , we see an enhancement in both analyses.

- In INCLUSIVE analysis we obtain

$M(h_c) = 3524.8 \pm 0.7(\text{stat}) \pm \sim 1(\text{syst}) \text{ MeV}$,

$B(\psi' \rightarrow \pi^0 h_c) \times B(h_c \rightarrow \gamma \eta_c) = (2-6) \times 10^{-4}$,

significance $> 3 \sigma$.

Thus,

$\Delta M_{hf} \equiv \langle M(\chi_J) \rangle - M(^1P_1) = 0.5 \pm 0.7(\text{stat}) \pm \sim 1(\text{syst}) \text{ MeV}$

- In EXCLUSIVE analysis we obtain

$M(h_c) = 3524.4 \pm 0.9(\text{stat}) \text{ MeV}$,

significance $\sim 5 \sigma$.

- The inclusive and exclusive results for $M(h_c)$ are in excellent agreement.