7 Overall summary

In the previous sections we have outlined in detail both the design and physics goals of the proposed CLEO III detector. A separate document summarizes this proposal and provides the detailed budgets, schedules, and construction time lines. Although this document has contributions from a large number of physicists in the CLEO collaboration, the formulation, direction, and overall coordination was carried out by the Inner Detector Task Force of C. Bebek, H. Kagan (Chairperson), R. Kass, H. Nelson, E. Nordberg, M. Ogg, D. Peterson, D. Rice, S. Stone, and M. Witherell.

References

[1] R. Ammar *et al.*, Phys. Rev. Lett. **71**, 674 (1993).

[2] M. Battle *et al.*, "Observation of B^0 Decay to Two Charmless Mesons," CLNS 93/1235 (1993).

[3] M. Gronau and D. Wyler, Phys. Lett. **265B**, 172 (1991).

[4] L. Wolfenstein, Phys. Rev. **D31**, 2381 (1985)

[5] B. Stech, Mod. Phys. Lett. **A6**, 3113 (1991); M. Neubert and B. Stech, Phys. Rev. **D44**, 775 (1991); K. Maltman *et al.*, Nucl. Phys. **A508**, 371c (1990); W. A. Bardeen, A. J. Buras and J. M. Gerard, Phys. Lett. **192B**, 138 (1987); M. B. Gavela *et al.*, Phys. Lett. **154B**, 425 (1985); and references contained therein.

[6] N. Isgur and M. Wise, Phys. Lett. **237B**, 527 (1990).

[7] N. Isgur, D. Scora, B. Grinstein and M. Wise, Phys. Rev. **D39**, 799 (1989).

[8] J. G. Korner and G. A. Stech, Zeit. Phys. **C38**, 511 (1988) and Zeit. Phys. **C41**, 690 (1988).

[9] M. Bauer, B. Stech and M. Wirbel, Zeit. Phys. **C29**, 637 (1985).

[10] M. Neubert, "Model-Independent Extraction of V_{cb} from Semi-Leptonic Decays," Heidelberg preprint HD-THEP-91-13 (1991).

[11] J. Bartelt *et al.*, CLNS-93-1240, Sep. 1993.

[12] G. Altarelli *et al.*, Nucl. Phys. **B208**, 365 (1982).

[13] R. Fulton *et al.*, Phys. Rev. Lett. **64**, 16 (1990).

[14] H. Albrecht *et al.*, Phys. Lett. **234B**, 409 (1990).

[15] M. Pauli(ARGUS), "Proceedings of the Joint International Symposium and Europhysics Conference on High Energy Physics," ed. by S. Hegarty, K. Porter and E. Quercigh (World Scientific, Singapore, 1992), p. 592.

[16] J. D. Richman(CLEO), "Search for Exclusive $b\ell \to u$ Semileptonic Decays of B Mesons," Proc. of the 7th Meeting of the APS Division of Particles and Fields, FNAL, Nov. 10-14, 1992.

[17] For a review see N. G. Desphande, "Theory of Penguins in B Decays, in B **Decays**, ed. by S. Stone, World Scientific, Singapore, (1992), p430.

[18] J. Bartelt *et al.*, Phys. Rev. Lett. **71**, 4111 (1993).

[19] R. Grigjanis *et al.*, Phys. Lett. **237B**, 252 (1990); B. Grinstein, R. Springer and M. Wise, Phys. Lett. **202B**, 138 (1988).

[20] A. Ali and C. Greub, Phys. Lett. **259B**, 182 (1991).

[21] A. Ali, T. Ohl and T. Mannel "Radiative Rare B Decays into Higher K-Resonances," DESY 92-113 (1992); P. J. O'Donnell *et al.*, "Exclusive Rare Decay $B \to K^* \gamma$," UTPT-92-12 (1992); A. Ali and T. Mannel, Phys. Lett. **264B**, 447 (1991), erratum Phys. Lett. **274B**, 526 (1992); P. Colangelo *et al.*, Zeit. Phys. **C45**, 575 (1990); C. A. Dominguez, N. Paver and Riazuddin, Phys. Lett. **214B**, 459 (1988).

[22] N.G. Deshpande and P. O'Donnell, "Proceedings of the Workshop towards Establishing a b Factory," Syracuse, Sept., 1989, p. 1.35 and 1.66.

[23] D. Du and Z. Xing, Phys. Rev. **D48**, 4155 (1993).

[24] S. Weinberg, Phys. Rev. Lett. **37**, 657 (1976); G. Bélanger, C.Q. Geng, and P. Turcotte, Phys. Rev. **D48**, 5419 (1993); T. Hayashi, M. Matsuda, and M. Tanimoto, paper 305 submitted to the International Symposium on Lepton and Photon Interactions at High Energies, Cornell, 1993.

[25] E. H. Thorndike, Bull. Am. Phys. Soc. **38**, 922(1993); D. Besson, invited talk at "1993 International Symposium on Lepton and Photon Interactions at High Energies," Cornell.

[26] F. Borzumati and A. Masiero, Phys. Lett. **192B**, 437 (1987) and erratum Phys. Lett. **198B**, 590 (1987); R. Barbieri and G.F. Giudice, quoted by A. Sanda, International Symposium on Lepton and Photon Interactions at High Energies, Cornell, 1993.

[27] M. S. Alam *et al.*, Phys. Rev. Lett. **51**, 1143 (1983); M. S. Alam *et al.*, Phys. Rev. Lett. **59**, 22 (1987); H. Albrecht *et al.*, Zeit. Phys. **C42**, 519 (1989); G. Crawford *et al.*, Phys. Rev. **D45**, 752 (1992).

[28] C. Dominguez *et al.*, Phys. Lett. **197B**, 423 (1987); L. Reinders, Phys. Rev. **D38**, 947 (1988); estimates by Shifman and by Narison quoted by R. Schindler (SLAC-PUB-4995).

[29] C. Bernard *et al.*, Phys. Rev. **D38**, 3540 (1988).

[30] E. Eichten *et al.*, Phys. Rev. **D34**, 1547 (1986).

[31] I. Hinchliffe, preprint LBL-18742 (1984).

[32] R. Ammar *et al.*, "Search for B^0 decays to two charged leptons," CLNS 93/1258.

[33] Report of the Working Group on CP violation and rare decays, *Proceedings of the 1984 Snowmass Summer Study on the Design and Utilization of the Superconducting Super Collider.*

[34] S. Stone, "Possible Ways of Observing CP Violation at a Symmetric e^+e^- B Factory," Syracuse University preprint HEPSY 92-2 (1992); S. Stone, Nucl. Inst. Meth. **A333**, 15 (1993).

[35] We ignore the small contribution from color suppressed decays for diagram (a).

[36] For example, this would be true if only one amplitude dominates each of the decays to D^0 or $\bar{D}^0$.

[37] The phase δ in this discussion is the difference $\delta - \bar{\delta}$ between the hadronic phases δ and $\bar{\delta}$ of the two amplitudes A and Ā, respectively, that are defined by Gronau and Wyler.

[38] We use $\lambda \approx 0.22$ and a for the Wolfenstein parameter A to avoid confusion with our amplitude A.

[39] We use $|V_{ub}/V_{cb}| = 0.08 \pm 0.03$ and λ is taken as 0.22.

[40] P. H. Cox, S. Hovater and S. T. Jones, Phys. Rev. **D33**, 1503 (1986); J. H. K'uhn, S. Nussinov and R. R'uckl, Zeit. Phys. **C5**, 117 (1980).

[41] M. S. Alan *et al.*, CLNS 94-1270, submitted to Phys. Rev. D.

[42] K. Hikasa *et al.*, Phys. Rev **D45**, Part II(1992).

[43] This number is in agreement with that observed in the CLEO II data. It is possible to increase the efficiency by adding more D^0 decay modes and improving the detector solid angle for photons and the charged particle tracking efficiency. In all estimates we use 86% for the efficiency of reconstructing a high momentum track.

[44] G. Valencia, Phys. Rev. **D39**, 3339 (1992).

[45] A. Deandrea *et al.*, Phys. Lett. **318B**, 549 (1993); M. Bauer, B. Stech and M. Wirbel, Zeit. Phys. **C34**, 103 (1987).

[46] G. Kramer and R. Palmer, Phys. Rev. **D45**, 193 (1992); Phys. Lett. **279B**, 181 (1992).

[47] G. Burdman and J. F. Donoghue Phys. Rev. **D45**, 187 (1992).

[48] A. Bean(CLEO), "Measurement of $\mathcal{B}(D^+ \to \pi^0 \ell^+ \nu)/\mathcal{B}(D^+ \to \bar{K}^0 \ell^+ \nu)$," in proc. of the 7th Meeting of the APS Division of Particles and Fields, FNAL Nov. 10-14, 1992; private communication from Arne Freyberger.

[49] D. Bortoletto *et al.*, Phys. Rev. Lett. **69**, 2046 (1992).

[50] J. F. Amundson *et al.*, Phys. Rev. **D47**, 3059 (1003).

[51] D. Acosta *et al.*, CLNS 93-14 (1993).

[52] F. Muheim and S. Stone, "Three Model Dependent Estimates of $\mathcal{B}(D_s^+ \to \phi\pi^+)$," HEPSY 93-3 (1993), to be published in Phys. Rev. D.

[53] I. Bigi, Proceedings of the Charm Physics Workshop, Peking(1987); I. Bigi and A. I. Sanda, Phys. Lett. **171B**, 320 (1986); J. Donoghue, E. Golowich, B. R. Holstein and J. Trampetic, Phys. Rev. **D33**, 179 (1986); L. Wolfenstein, Phys. Lett. **164B**, 170 (1985); P. Colangelo and G. Nardulli, Phys. Lett. **242B**, 71 (1990); H. Georgi, Phys. Lett. **297B**, 353 (1992); T. Ohi, G. Ricciardi and E. H. Simmons, Nucl. Phys. **B403**, 605 (1993).

[54] D. S. Akerib *et al.*(CLEO), "Observation of $D^0 \to K^+\pi^-$," presented at Photon Lepton Conference, Ithaca NY(1993).

Singapore (1992).

[55] J. Haissinski, Orsay Preprint No. LAL-90-64, 1990, to be published in Proceedings of the 8th EPS General Conference: Trends in Physics (Amsterdam, The Netherlands, 1990).

[56] S. L. Glashow, A. Salam and S. Weinberg, Rev. Mod. Phys. **52**, 515 (1980).

[57] For a recent review, see for example, K. K. Gan and M. L. Perl, Int. J. Mod. Phys. **A3**, 531 (1988); B. C. Barish and R. Stroynowski, Phys. Rep. **157**, 1 (1988).

[58] See for example, P. Langacker, B. Sathiapalan and G. Steigman, Nucl. Phys. **B266**, 669 (1986) and references therein.

[59] H. Albrecht *et al.*, Phys. Lett. **202B**, 149 (1988).

[60] Y. S. Tsai, Phys. Rev. **D4**, 2821 (1971).

[61] F. J. Gilman and D. H. Miller, Phys. Rev. **D17**, 1846 (1978).

[62] H. B. Thacker and J. J. Sakurai, Phys. Lett. **36B**, 103 (1971).

[63] D. Akerib *et al.*, Phys. Rev. Lett. **69**, 3610 (1992), *ibid.* Phys. Rev. Lett. **71**, 3395 (1993).

[64] T. Das, V. S. Mathur and S. Okubo, Phys. Rev. Lett. **18**, 761 (1967).

[65] M. Artuso *et al.*, Phys. Rev. Lett. **69**, 3278 (1992).

[66] F. J. Gilman, Phys. Rev. **D35**, 3541 (1987).

[67] A. Pich, Phys. Lett. **196B**, 561 (1987).

[68] C. Leroy and J. Pestieau, Phys. Lett. **72B**, 398 (1978); S. N. Biswas *et al.*, Phys. Lett. **80B**, 393 (1979); N. Paver and D. Treleani, Lett. Nuovo Cimento **31**, 364 (1981); V. P. Barannik, A. P. Korzh and M. P. Rekalo, Acta Phys. Pol. **B13**, 835 (1982).

[69] See for example, H. Harari, Fundamental Forces, edited by D. Frame and K. J. Peach (St. Andrews, 1984), p.357; R. D. Pecci, New and Exotic Phenomena, edited by O. Fackler and J. Tran Thanh Van (Les Arcs, Savoie, France, 1987), p.431.

[70] A. Bean *et al.*, CLEO preprint, CLNS 92/1172, submitted to Phys. Rev. Lett.

[71] H. Albrecht *et al.*, Phys. Lett. **246B**, 278 (1990).

[72] W. Fetscher, H.-J. Gerber and K. F. Johnson, Phys. Lett. **173B**, 102 (1986).

[73] W. Fetscher, in Proposal for an Electron Positron Collider for Heavy Flavour Particle Physics and Synchrotron Radiation, PSI PR-88-09 (1988).

[74] M. Artuso *et al.*, "Measurement of the Cross-Section for $\gamma\gamma \rightarrow p\bar{p}$," CLNS93/1245, submitted to Phys. Rev. D.

[75] D. Acosta *et al.*, "Two-Photon Production of Charged Pion and Kaon Pairs," in preparation for Phys. Rev. D.

[76] V. Shelkov *et al.*, "Measurement of the Two-Photon Production of the χ_{c2}," CLNS93/1250, accepted by Phys. Rev. D.

[77] R. Sonnenblick *et al.*, Nucl. Inst. Meth. **A310**, 189 (1991).

[78] G. Hall *et al.*, Nucl. Inst. Meth. **A326**, 228 (1993).

[79] J. F. Bak *et al.*, Nucl. Phys. **B288**, 681 (1987).

[80] Private communication, members of the L3 collaboration.

[81] Note that Radeka (*Low-Noise Techniques in Detectors*, Ann. Rev. Nucl. Part. Sci, **38** (1988) 217) discusses series noise in a different limit, namely where RC is large; in this limit, $ENC^2 \sim 1/\tau$. In our case, however, RC is small, and the resulting ENC is independent of the integration time.

[82] W. Trischuk, private communication. See also comments in O. Toker *et al.*, "Viking, a CMOS low noise monolithic 128 channel frontend for Si-strip detector readout," CERN-PPE/93-141.

[83] J. P. Alexander *et al.*, *Studies of Double-Sided, Double Metal Silicon Strip Detectors*, contribution to the International Symposium on the Development and Application of Semiconductor Tracking Detectors, to appear in Nucl. Instr. Meth..

[84] Current versions of the VIKING chip yield a noise slope as low as 12.5/pF, with a 2.0 μs shaping time. A new version, now being tested, appears to offer 5.4/pF (private communication, P. Weilhammer.) We are presently exploring the possibilities of using such a chip, suitably tailored to CLEO needs.

[85] Such kapton circuits have been fabricated for gas microstrip detectors by CLEO collaborators at Ohio State University.

[86] W. Snoeys *et al.*, Nucl. Inst. Meth. **A326**, 144 (1993); See also, S. Parker, Nucl. Inst. Meth. **A275**, 494 (1989).

[87] *cf.* for instance J. C. Jernigan, SLAC-PUB-5925.

[88] D. G. Cassel *et al.*, Nucl. Inst. Meth. **A52**, 325 (1986).

[89] W. O. Miller *et al.*, Los Alamos Nat. Lab. report LA-12029.

[90] D. Hale, Univ. of California, Santa Barbara, private communication.

[91] W. Adam *et al.*, "The Ring Imaging Cherenkov Detector of DELPHI," presented at First Int. Conf. on RICH Detectors, Bari, Italy, (1993); K. Abe *et al.*, "Performance of the CRID at SLD," ibid.; U. M'uller *et al.*, "Particle Identification with the RICH Detector in Experiment WA89 at CERN," ibid.; H.-W. Siebert *et al.*, "The OMEGA-RICH," ibid.; R. Baur *et al.*, "The CERES RICH Detector System," ibid.; A. Braem *et al.*, "Fast Rich Detector with a CsI Photocathode at Atmospheric Pressure," ibid.; M. Staric *et al.*, "Measurement of Cherenkov Angles with a CsI Photocathode in a Multiwire Chamber," ibid.

[92] M. French *et al.*, Nucl. Inst. Meth. **A324**, 511 (1990).

[93] R. Arnold *et al.*, Nucl. Inst. Meth. **A314**, 465 (1992).

[94] J. Seguinot *et al.*, Nucl. Inst. Meth. **A297**, 133 (1990).

[95] For more details see M. Artuso and Y. Mukhin, "Comparison of Expected Performance of Fast Ring Imaging Cherenkov Counter Systems," CBX 93-45 (1993); M. Artuso and Y. Mukhin, "A Fast Ring Imaging Detector for the CLEO Upgrade," presented at First Int. Conf.on RICH Detectors, Bari, Italy, (1993), to appear in Proceedings.

[96] D. Anderson *et al.*, Nucl. Inst. Meth. **A323**, 626 (1992); J. Seguinot and T. Ypsilantis, Ring Imaging Cherenkov Counters, CERN Preprint CERN–LAA/PI/91-004 (1991); D. F. Anderson, S. Kwan and V. Peskov, "CsI and Some New Photocathodes," FERMILAB-Conf-93/128 (1993); R. Aleksan *et al.*, "Measurements of CsI Photocathode quantum efficiency in Methane," presented at First Int. Conf. on RICH Detectors, Bari, Italy, (1993).

[97] S. Ilie and G. Lenzen, "Perfluorocarbon Liquid, Specific Chemical Aspects For Use in the DELPHI RICH," DELPHI 93-33 (1993).

[98] R. Brandelik *et al.*, Phys. Lett. **83B**, 261 (1979); A. Onuchin *et al.*, Nucl. Inst. Meth. **A315**, 517 (1992).

[99] H. Burkhardt *et al.*, Nucl. Inst. Meth. **184**, 319 (1981); M. Bockhorst, "Measurement of the optical properties of the components of an aerogel Cerenkov counter with wavelength shifter light guide," BONN-IR-92-35, Sep 1992.

[100] G. Eigen, D. G. Hitlin, M. Lakata, J. Y. Oyang, and W. J. Wisniewski, "Silica aerogel Cerenkov detectors for particle identification," SLAC-BABAR-NOTE-100(1992); G. Eigen, D.G. Hitlin and J.Y. Oyang, "Studies of silica aerogel as a low mass particle identification system for the SLAC B factory detector," SLAC-BABAR-NOTE-101(1993).

[101] M. Selen and K. Honscheid, *Particle ID via Cherenkov Correlated Timing*, CLEO Preprint CBX 92-116; see also paper of same name by K. Honscheid, M. Selen and M. Sivertz, Proceedings of the First Int. Conf. on RICH Detectors, Bari, Italy, 1993.

[102] The idea of using total internal reflection to guide the Cherenkov light and to measure the position and shape of the Cherenkov rings as projected from the end of the radiator was first discussed by Blair Ratcliff. For a review of his ideas, refer to his talk "Status of the B Factory Detector for PEP-II: A Progress Report" at the XXVI. International Conference on High Energy Physics, Dallas (1992) (SLAC-PUB 5946 (92)). The first usage of total-internal reflection in Cherenkov counters was suggested by Fitch and Motley (Phys. Rev. **101**, 496 (1956)).

[103] This is a conservative assumption. The fastest photo-tube available from Hammamatsu has a single electron transit time jitter of 160 ps FWHM.

[104] Advanced Photonics Inc., Camarillo, California 93012.

[105] M. Grande and G. R. Moss, Nucl. Inst. Meth. **215**, 539 (1983).

[106] T. Browder *et al.*, "CESR-B R&D Report, June 1993," Cornell University, Ithaca, N.Y. page 4-1.