

Advanced Accelerator Physics and Accelerator Simulation Homework 6

Georg H. Hoffstaetter
Physics 488/688, Cornell University

March 9, 2007

Exercise 1:

Given a reference trajectory that is a helix around the z -axis with

$$\vec{R}(z) = r \cos(kz) \vec{e}_X + r \sin(kz) \vec{e}_Y + z \vec{e}_Z, \quad (1)$$

with the Cartesian coordinate vectors \vec{e}_X , \vec{e}_Y and \vec{e}_Z .

(a) Show that z is not the pathlength s with which the reference trajectory is parametrized. Then compute the path length $s(z)$ and specify $\vec{R}(s)$ so that $|d\vec{R}| = ds$ and compute \vec{e}_s , \vec{e}_κ , and \vec{e}_b .

(b) Compute \vec{e}_x and \vec{e}_y of the curvilinear system and check that $\frac{d}{ds}\vec{e}_x$ and $\frac{d}{ds}\vec{e}_y$ are what they are specified to be in the handouts.

Exercise 2:

Let the linearized particle transport from initial phase space coordinates \vec{z}_i to final phase space coordinates \vec{z}_f be:

$$\begin{pmatrix} x_f \\ a_f \\ \tau_f \\ \delta_f \end{pmatrix} = \begin{pmatrix} M_{11} & M_{12} & 0 & D_x \\ M_{21} & M_{12} & 0 & D_a \\ T_x & T_a & 1 & R_{56} \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_i \\ a_i \\ \tau_i \\ \delta_i \end{pmatrix}. \quad (2)$$

The zeros in the matrix show that the particle motion is independent of the starting time and that the energy is independent of the starting conditions.

(a) Describe the meaning of the coefficients D_x , D_a , T_x , and T_a .

(b) Use the fact that this 4×4 matrix is symplectic to show that the top left-hand 2×2 sub-matrix is symplectic.

(c) Show how T_x and T_a can be computed when this top left-hand sub-matrix and the dispersion D_x and its slope D_a are known.

Exercise 3:

Using the relativistic approximation $\gamma^2 \gg 1$, derive the 6 dimensional transport matrix for

A) a quadrupole magnet with length L and quadrupole coefficient k . Make a separate calculation for positive and for negative k .

B) a combined function magnet with length L , curvature κ and quadrupole coefficient k . Use $K = k + \kappa^2$ and distinguish the cases $k > 0$, $k < -\kappa^2$, and $-\kappa^2 < k < 0$.

C) Check that the time of flight could have been computed from the condition $\vec{T} = -\mathbf{J}_4 \mathbf{M}_4^{-1} \vec{D}$ due to symplecticity.

Exercise 4:

A time of flight spectrometer takes all particles that come from a collision point regardless of their initial slopes x' and y' and transports them to a point in a detector plane. The time of flight should depend only on the energy, not on the initial position or the initial angle of the particles in the collision plane. Write the most general form that the transport matrix from the collision plane to the detector plane can have.

Exercise 5:

If not the time of flight $\tau = (t_0 - t) \frac{E_0}{P_0}$ and the relative energy change $\delta = \frac{\Delta E}{E}$ had been chosen as phase space variables, but the deviation in path length Δl and the relative momentum deviation $\frac{\Delta P}{P}$, how would the transport matrix look like and how could it be computed from the transport matrix in exercise 4?