



Time of Flight from Symplecticity



CHESS & LEPP

$$\underline{M} = \begin{pmatrix} \underline{M}_4 & \vec{0} & \vec{D} \\ \vec{T}^T & 1 & M_{56} \\ \vec{0}^T & 0 & 1 \end{pmatrix} \text{ is in SU(6) and therefore } \underline{M} \underline{J} \underline{M}^T = \underline{J}$$

$$\begin{pmatrix} \underline{M}_4 \underline{J}_4 & -\vec{D} & \vec{0} \\ \vec{T}^T \underline{J}_4 & -M_{56} & 1 \\ \vec{0}^T & -1 & 0 \end{pmatrix} \begin{pmatrix} \underline{M}_4^T & \vec{T} & \vec{0} \\ \vec{0}^T & 1 & 0 \\ \vec{D}^T & M_{56} & 1 \end{pmatrix} = \begin{pmatrix} \underline{J}_4 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} \underline{M}_4 \underline{J}_4 \underline{M}_4^T & \underline{M}_4 \underline{J}_4 \vec{T} - \vec{D} & \vec{0} \\ \vec{T}^T \underline{J}_4 \underline{M}_4^T + \vec{D}^T & 0 & 1 \\ \vec{0}^T & -1 & 0 \end{pmatrix} = \begin{pmatrix} \underline{J}_4 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$$

$$\vec{T} = -\underline{J}_4 \underline{M}_4^{-1} \vec{D}$$

It is sufficient to compute the 4D map \underline{M}_4 , the Dispersion \vec{D} and the time of flight term M_{56}



$$x'' = -x k$$

$$y'' = y k$$

$$\underline{M}_4 = \begin{pmatrix} \cos(\sqrt{k} s) & \frac{1}{\sqrt{k}} \sin(\sqrt{k} s) & & \\ -\sqrt{k} \sin(\sqrt{k} s) & \cos(\sqrt{k} s) & & \\ & & \underline{0} & \\ & & & \cosh(\sqrt{k} s) & \frac{1}{\sqrt{k}} \sinh(\sqrt{k} s) \\ & & & \sqrt{k} \sinh(\sqrt{k} s) & \cosh(\sqrt{k} s) \end{pmatrix}$$

As for a drift:

$$\vec{D} = \vec{0} \Rightarrow \vec{T} = \vec{0}$$

$$M_{56} = 0$$

For $k < 0$ one has to take into account that

$$\cos(\sqrt{k} s) = \cosh(\sqrt{|k|} s), \quad \sin(\sqrt{k} s) = i \sinh(\sqrt{|k|} s)$$

$$\cosh(\sqrt{k} s) = \cos(\sqrt{|k|} s), \quad \sinh(\sqrt{k} s) = i \sin(\sqrt{|k|} s)$$



The Combined Function Bend



CHES & LEPP

$$x'' = -x \underbrace{(\kappa^2 + k)}_K + \delta \kappa$$

$$y'' = y k, \quad \tau' = -\kappa x$$

$$\underline{M}_6 = \begin{pmatrix} \underline{M}_x & \underline{0} & \vec{0} \vec{D} \\ \underline{0} & \underline{M}_y & \underline{0} \\ \underline{T} & \underline{0} & \underline{M}_\tau \end{pmatrix}$$

$$\underline{M}_x = \begin{pmatrix} \cos(\sqrt{K} s) & \frac{1}{\sqrt{K}} \sin(\sqrt{K} s) \\ -\sqrt{K} \sin(\sqrt{K} s) & \cos(\sqrt{K} s) \end{pmatrix}$$

$$\underline{M}_y = \begin{pmatrix} \cosh(\sqrt{k} s) & \frac{1}{\sqrt{k}} \sinh(\sqrt{k} s) \\ \sqrt{k} \sinh(\sqrt{k} s) & \cosh(\sqrt{k} s) \end{pmatrix}$$

$$\vec{D} = \begin{pmatrix} \frac{\kappa}{K} [1 - \cos(\sqrt{K} s)] \\ \frac{\kappa}{\sqrt{K}} \sin(\sqrt{K} s) \end{pmatrix}$$

$$\underline{M}_\tau = \begin{pmatrix} 1 & M_{56} \\ 0 & 1 \end{pmatrix}$$

$$M_{56} = \frac{\kappa^2}{K\sqrt{K}} [\sin(\sqrt{K} s) - \sqrt{K} s]$$

\underline{T} from symplecticity

Options:


- For $k > 0$:
focusing in x, defocusing in y.
- For $k < 0, K < 0$:
defocusing in x, focusing in y.
- For $k < 0, K > 0$:
weak focusing in both planes.



The Thin Lens Approximation



CHESS & LEPP



$$\vec{z}(s) = \underline{M}(s) \vec{z}_0 = \underline{D}\left(\frac{s}{2}\right) \underline{D}^{-1}\left(\frac{s}{2}\right) \underline{M}(s) \underline{D}^{-1}\left(\frac{s}{2}\right) \underline{D}\left(\frac{s}{2}\right) \vec{z}_0$$


Drift: $\underline{M}_{\text{drift}}^{\text{thin}}(s) = \underline{D}^{-1}\left(\frac{s}{2}\right) \underline{M}(s) \underline{D}^{-1}\left(\frac{s}{2}\right) = \underline{1}$



The Thin Lens Quadrupole



CHESS & LEPP



$$\begin{aligned} \underline{M}_{\text{quad},x}^{\text{thin}}(s) &= \begin{pmatrix} 1 & -\frac{s}{2} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos(\sqrt{k}s) & \frac{1}{\sqrt{k}} \sin(\sqrt{k}s) \\ -\sqrt{k} \sin(\sqrt{k}s) & \cos(\sqrt{k}s) \end{pmatrix} \begin{pmatrix} 1 & -\frac{s}{2} \\ 0 & 1 \end{pmatrix} \\ &\approx \begin{pmatrix} 1 & -\frac{s}{2} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & s \\ -ks & 1 \end{pmatrix} \begin{pmatrix} 1 & -\frac{s}{2} \\ 0 & 1 \end{pmatrix} \approx \begin{pmatrix} 1 & -\frac{s}{2} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \frac{s}{2} \\ -ks & 1 + \frac{ks^2}{2} \end{pmatrix} \end{aligned}$$

Weak magnet limit: $\sqrt{k}s \ll 1$

$$\underline{M}_{\text{quad},x}^{\text{thin}}(s) \approx \begin{pmatrix} 1 & 0 \\ -ks & 1 \end{pmatrix}$$