

---

# Introduction to the SM (5)

Yuval Grossman

Cornell

# Yesterday...

---

- Yesterday: Symmetries
- Today
  - SSB
  - the SM

---

# SSB

# Breaking a symmetry

---



# SSB

---

- By choosing a ground state we break the symmetry
- We choose to expand around a point that does not respect the symmetry
- PT only works when we expand around a minimum

What is the different between a broken symmetry and no symmetry?

SSB implies relations between parameters

# SSB

---

Symmetry is  $x \rightarrow -x$  and we keep up to  $x^4$

$$f(x) = a^2 x^4 - 2b^2 x^2 \quad x_{\min} = \pm b/a$$

We choose to expand around  $+b/a$  and use  $u \rightarrow x - b/a$

$$f(x) = 4b^2 u^2 + 4bau^3 + a^2 u^4$$

- No  $u \rightarrow -u$  symmetry
- The  $x \rightarrow -x$  symmetry is hidden
- A general function has 3 parameters  $c_2 u^2 + c_3 u^3 + c_4 u^4$
- SSB implies a relation between them

$$c_3^2 = 4c_2 c_4$$

# SSB in QFT

---

- When we expand the field around a minimum that is not invariant under a symmetry

$$\phi \rightarrow v + h$$

- It breaks the symmetries that  $\phi$  is not a singlet under
- Masses to other fields via Yukawa interactions

$$\phi X^2 \rightarrow (v + h)X^2 = vX^2 + \dots$$

- Gauge fields of the broken symmetries also get mass

$$|D_\mu \phi|^2 = |\partial_\mu \phi + iqA_\mu \phi|^2 \ni A^2 \phi^2 \rightarrow v^2 A^2$$

---

# The SM

# The SM

---

Input: Symmetries and fields

- Symmetry: 4d Poincare and

$$SU(3)_C \times SU(2)_L \times U(1)_Y$$

- Fields:

- 3 copies of QUDLE fermions

$$\begin{array}{lll} Q_L(3, 2)_{1/6} & U_R(3, 1)_{2/3} & D_R(3, 1)_{-1/3} \\ L_L(1, 2)_{-1/2} & E_R(1, 1)_{-1} & \end{array}$$

- One scalar

$$\phi(1, 2)_{+1/2}$$

# Then Nature is described by

---

- Output: the most general  $\mathcal{L}$  up to dim 4

$$\mathcal{L} = \mathcal{L}_{kin} + \mathcal{L}_{Higgs} + \mathcal{L}_{Yukawa}$$

- This model has a  $U(1)_B \times U(1)_e \times U(1)_\mu \times U(1)_\tau$  accidental symmetry
- Initial set of measurements to find the parameters
  - SSB:  $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$
  - Fermion masses, gauge couplings and mixing angles

The SM pass (almost) all of its tests

---

# The gauge interactions

# The gauge part

---

$$SU(3)_C \times SU(2)_L \times U(1)_Y \rightarrow SU(3)_C \times U(1)_{EM}$$

Three parts, each look so different...

- QED - photon interaction: Perturbation theory
- QCD - gluon interaction: Confinement and asymptotic freedom
- Electro-weak: SSB and massive gauge bosons

# $\mathcal{L}_{kin}$ and $SU(2) \times U(1)$

---

- Four gauge bosons DOFs

$$W_a^\mu(1, 3)_0 \quad B^\mu(1, 1)_0$$

- The covariant derivative is

$$D^\mu = \partial^\mu + igW_a^\mu T_a + ig'Y B^\mu$$

- Two parameters  $g$  and  $g'$
- $Y$  is the  $U(1)$  charge of the field  $D_\mu$  work on
- $T_a$  is the  $SU(2)$  representation
- $T_a = 0$  for singlets.  $T_a = \sigma_a/2$  for doublets
- Write  $D_\mu$  for  $L(1, 2)_{-1/2}$  and  $E(1, 1)_{-1}$

# Explicit examples

---

$$D^\mu = \partial^\mu + igW_a^\mu T_a + ig'Y B^\mu$$

- Write  $D_\mu$  for  $L(1, 2)_{-1/2}$  and  $E(1, 1)_{-1}$

$$D^\mu L = \left( \partial^\mu + \frac{i}{2}gW_a^\mu \sigma_a - \frac{i}{2}g' B^\mu \right) L$$

$$D^\mu E = (\partial^\mu - ig' B^\mu) E$$

- HW: Using  $\phi(1, 2)_{1/2}$  write  $D^\mu \phi$

# QED

---

- Where is QED in all of this?

$$Q = T_3 + Y$$

- We can write explicitly for  $L(1, 2)_{-1/2}$  and  $\phi(1, 2)_{1/2}$

$$L_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \quad \phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$$

- This is arbitrary. It becomes useful once we have SSB

# SSB in the SM

---

$$-\mathcal{L}_{Higgs} = \lambda\phi^4 - \mu^2\phi^2 = \lambda(\phi^2 - v^2)^2$$

- We measure the fact that  $\mu^2 > 0$  by having SSB
- The minimum is at  $|\phi| = v$
- $\phi$  has 4 DOFs. We can choose

$$\langle\phi_1\rangle = \langle\phi_2\rangle = \langle\phi_4\rangle = 0 \quad \langle\phi_3\rangle = v$$

- It leads to:  $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$
- We call the remaining symmetry EM
- Could we “choose” the vev in the neutral direction?
- We left with one real scalal field: the Higgs boson

---

# Spectrum

# Gauge boson masses

---

- $W_1, W_2, W_3, B$
- Gauge bosons masses from  $|D_\mu\phi|^2$  (HW: do it)
- Diagonalizing the mass matrix the masses are

$$M_{W^+}^2 = M_{W^-}^2 = \frac{1}{4}g^2v^2 \quad M_Z^2 = \frac{1}{4}(g^2 + g'^2)v^2 \quad M_A^2 = 0$$

- The mass eigenstates

$$W^\pm = \frac{1}{\sqrt{2}}(W_1 \mp iW_2) \quad \tan\theta_W \equiv \frac{g'}{g}$$

$$Z = \cos\theta_W W_3 - \sin\theta_W B \quad A = \sin\theta_W W_3 + \cos\theta_W B$$

- We have a  $\theta_W$  rotation from  $(W_3, B)$  to  $(Z, A)$

# $\mathcal{L}_{Yuk}$ and fermion masses

---

- There is no way to write a mass term (why?)
- The Yukawa part of the leptons

$$\mathcal{L}_{Yuk} = y_{ij} \overline{L}_{Li} E_{Rj} \phi \Rightarrow m_{ij} \overline{L}_{Li} E_{Rj} \quad m_{ij} = v y_{ij}$$

- $i, j = 1, 2, 3$  are flavor indices
- $y$  is a general complex  $3 \times 3$  matrix and we can choose a basis where  $m$  is diagonal and real

$$m_{ij} = y v = \text{diag}(m_e, m_\mu, m_\tau)$$

- Neutrinos are massless

---


# Interactions

# Charged current interactions

$$-\frac{g}{\sqrt{2}} \bar{\nu}_{eL} W^\mu \gamma_\mu e_L^- + h.c.$$

- Only left-handed fields take part in charged-current interactions. Therefore the  $W$  interaction violate parity
- The  $W\ell\nu$  interaction is universal
- Can be used to measure  $g$

$$A \sim g^2/m_W^2 \sim G_F$$



# Neutral currents

---

$$\mathcal{L}_{\text{int}} = \frac{e}{\sin \theta \cos \theta} (T_3 - \sin^2 \theta_W Q) \bar{\psi} \gamma^\mu \psi Z_\mu + e Q \bar{\psi} \gamma^\mu \psi A_\mu,$$

- We define

$$Q = T_3 + Y \quad e = g \sin \theta$$

- Photon coupling is parity invariant
- $Z$  couples to both LH and RH fermions but in a parity violating way
- The coupling to the  $Z$  is larger. So why we call it weak interaction?
- Once we know  $e$  and  $g$  we know  $\theta$

# The $\rho = 1$ relation

---

We get the following testable relation

$$\rho \equiv \frac{M_W^2}{M_Z^2 \cos^2 \theta_W} = 1 \quad \tan \theta_W \equiv \frac{g'}{g}$$

**The above is a signal of SSB**

# Experimental tests

---

- High energy: Open your pdg and check  $W$  and  $Z$  decays to leptons. What do you expect to see?
- $Z$  decays to lepton actually measures  $\sin^2 \theta_W \approx 0.23$
- HW: Calculate  $\Gamma(Z \rightarrow \nu\bar{\nu})/\Gamma(Z \rightarrow e^+e^-)$ , get  $\sin^2 \theta_W$  from the data and check the  $\rho = 1$  prediction
- Also low energy data tests

# Experimental tests of $\rho = 1$

- From the  $\rho = 1$  relation

$$\frac{m_W^2}{m_Z^2} = \cos^2 \theta_W \approx \left(\frac{80.4}{91.2}\right)^2 \approx 0.77 \Rightarrow \sin^2 \theta_W \approx 0.23$$

- $Z$  decays to leptons

$$\Gamma(Z \rightarrow \ell\ell) \sim \sum_{L,R} (T_3 - Q \sin^2 \theta_W)^2$$

$$\Gamma(Z \rightarrow \ell^+ \ell^-) \sim (1/2 - \sin^2 \theta_W)^2 + (\sin^2 \theta_W)^2 \sim 1/8$$

$$\Gamma(Z \rightarrow \nu\bar{\nu}) \sim (1/2)^2 \sim 1/4 \Rightarrow r \equiv \Gamma_\ell / \Gamma_{\text{inv}} \sim 1/6$$

- PDG:**  $\Gamma_\ell = 3.37\%$  and  $\Gamma_{\text{inv}} = 20.00\%$   $\Rightarrow r \sim 1/6$